

CLUBROOT SOIL TEST

1. Choose a sample point in a field entrance or highly trafficked area.
2. Clear loose organic matter from the topsoil.
3. Collect 2 cups of topsoil.
 - A composite sample may be composed by evenly mixing soil taken from each sampling point.
4. Let sample **air dry** before placing in a plastic, resealable bag. A cooler is not required.
5. Ship or Drop Off the soil sample to;
507-11 Ave Nisku, Alberta T9E 7N5

SEND IN A SAMPLE TODAY

LEARN MORE AT 2020SEEDLABS.CA

1-877-420-2099

20  20 Seed Labs Inc.
never stop growing

CLUBROOT SOIL TEST

- The best time to sample soil is spring and late fall
- 20/20 Seed Labs offers qualitative testing on ½ C soil or semi-quantitative testing on 5 g samples of soil. These are the highest amounts of soil tested in the industry using DNA techniques.
- **DO NOT** test a soil sample from fields that have a susceptible host plant and do not sample soil near a host plant. Test infected plants instead.
- **DO NOT** sample randomly across the field. Disease incidence decreases as you move into the field.
- A **positive test result** indicates that under the right environmental conditions and in a susceptible host, there is a potential for the disease to occur.

CLUBROOT DISEASE & MANAGEMENT



OPEN YOUR PHONE CAMERA & HOLD YOUR
FINGER ON THE CODE TO GET THE LINK
WWW.CANOLACOUNCIL.ORG